

Off-Diagonal Long-Range Order in Solid ^4He

Bryan K. Clark¹ and D. M. Ceperley²

¹*Dept. of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA*

²*NCSA and Dept. of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA*

Measurements of the moment of inertia by Kim and Chan have found that solid ^4He acts like a supersolid at low temperatures. To understand the order in solid ^4He , we have used Path Integral Monte Carlo to calculate the off-diagonal long range order (ODLRO) (equivalent to Bose-Einstein condensation, BEC). We do not find ODLRO in a perfect hcp crystal at the melting density. We discuss these results in relation to proposed quantum solid trial functions. We conclude that the solid ^4He wave function has correlations which suppress both vacancies and BEC.

PACS numbers: PACS: 67.80.-s, 67.40.-w, 02.70.Ss

Whether bulk solid has a region of its equilibrium phase diagram with supersolid behavior has been of theoretical interest[1, 2, 3] for many years. Recently, Kim and Chan[4] have observed behavior similar to that of non-classical rotational inertia (NCRI) at temperatures below 0.2K. Analogous to behavior in a superfluid, NCRI would be expected in a supersolid, a state of matter in which crystallinity and superfluid behavior coexist. Two key theoretical quantities in establishing the order of a bosonic system are the superfluid fraction (estimated[5] to vanish) and off-diagonal long range order (ODLRO) previously estimated only by variational methods.

Off-diagonal order (equivalent to a non-zero condensate fraction) tells us whether atoms at one end of the solid are in phase with atoms at the other end of the solid and would supply a mechanism for NCRI: for example, superfluid ^4He in three dimensions has both ODLRO and NCRI. ODLRO[6], is a property of the single particle density matrix:

$$n(\mathbf{r}, \mathbf{r}'; \beta) = \frac{V}{Z} \sum_{\alpha} \int d\mathbf{r}_2 \dots d\mathbf{r}_n e^{-\beta E_{\alpha}} \times \Psi_{\alpha}(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_n) \Psi_{\alpha}^*(\mathbf{r}', \mathbf{r}_2, \dots, \mathbf{r}_n) \quad (1)$$

where Ψ_{α} and E_{α} are the α^{th} eigenfunction/value of the many body Hamiltonian, Z the partition function, V the volume and $\beta = 1/k_b T$. A system is said to be Bose-condensed if an eigenvalue of the matrix $n(\mathbf{r}, \mathbf{r}')$ is proportional to the number of atoms in the system. In this paper, we consider a translationally invariant system in periodic boundary conditions. This implies that $n(\mathbf{r}, \mathbf{r}')$ only depends on $\Delta\mathbf{r} = \mathbf{r} - \mathbf{r}'$ and its eigenfunctions are momentum states. Considering the occupation of the $\mathbf{k} = 0$ momentum state, the condensate fraction is

$$n_0 = \frac{1}{V^2} \int d\mathbf{r} d\mathbf{r}' n(\mathbf{r}, \mathbf{r}'). \quad (2)$$

A system has ODLRO if the condensate fraction is greater than zero in the thermodynamic limit. That is: $n_0 = \lim_{|\Delta\mathbf{r}| \rightarrow \infty} n(\Delta\mathbf{r}) > 0$.

Reatto[8] showed that a pair-product (Jastrow) trial wave function (JWF), commonly used for liquid ^4He ,

has ODLRO. This trial function has the form $\Psi_J(R) = \exp(-\sum_{i<j} u(r_{ij}))$ where $u(r_{ij})$ is fixed by minimizing the variational estimate of the ground state energy. The Jastrow trial function is equivalent to classical system with a pair potential $v(r)$: the Jastrow factor maps into the classical pair potential ($v(r) \equiv 2k_B T u(r)$). The physical argument for BEC is straightforward: to calculate $n(\Delta\mathbf{r})$ one displaces an arbitrary atom by $\Delta\mathbf{r}$; if the average amplitude for this is non-zero, the system has BEC. The condensate fraction will be non-zero as long as $u(r)$ is short-ranged, except for special systems such as hard spheres at close packing density. It will also be true if the wave function has short-ranged three or higher body correlation factors. Chester[3] then pointed out that such a trial function can be a crystal, since it has the same distribution as a classical solid interacting with a pair potential. This argument can be used to prove[9] that there exist quantum systems with both crystalline order and BEC. Chester further conjectures that under the same conditions, a crystal will have BEC only if there exist ground state vacancies (or interstitials). This argument has been recently elaborated.[10]

The Jastrow form of trial function turns out to give a seriously incorrect density for freezing[11] and other properties. As is well-known, melting of classical systems occurs when the r.m.s. vibration around a lattice site equals about 0.14 of the nearest neighbor distance: Lindemann's melting criterion. However, solid ^4He does not melt until much later[12], with a Lindemann's ratio of 0.30. Assuming Lindemann's criteria for the melting of a classical system, no Jastrow trial function can describe a solid ^4He near melting, nor can it describe a quantum crystal without an intrinsic population of point defects.

A much better description of solid ^4He is obtained by multiplying the Jastrow function by a localization term; the Jastrow-Nosanow form[13] (JN) or "insulator" form:

$$\Psi_{JN}(R) = \frac{\Psi_J}{N!} \sum_P \prod_{i=1}^N \phi_{P_i}(\mathbf{r}_i). \quad (3)$$

Here $\phi_i(\mathbf{r})$ (a Wannier function) is localized about lattice

site i , and P a permutation of atoms to lattice sites. The sum over P projects out the bosonic component of the wave function; the projected wavefunction is a “permanent”. In fact, the symmetrization lowers the variational energy by an amount proportional to the exchange energy, calculated in ref. [25] to be about $3 \mu\text{K}/\text{atom}$; negligible for most properties but crucial for properties such as BEC or NCRI. The JN wave function is a very good description of the ground state judging by the computed energies, Debye-Waller factors, pair correlation function and estimated melting density[11]. However, the solid order is put in “by hand” rather than coming about spontaneously. Broken translational symmetry is the hallmark of the crystal state[15][16]. Upon freezing a density wave arises which defines the lattice sites, a mean field potential and hence $\phi_i(\mathbf{r})$. It has been shown[14] that this type of trial function does not have BEC, assuming that the sum of the overlap of ϕ with those on other sites is less than unity as is expected to be the case. Because of the localized functions, the Reatto-Chester theorem on BEC does not apply.

An alternative trial function[17, 18, 19] (we will denote as a metal, Ψ_M) is obtained by making the single-body function $\phi(\mathbf{r})$ independent of the lattice site, but still having lattice symmetry. For electrons, $\phi(\mathbf{r})$ would be a Bloch function of band theory and would be obtained as a solution of a mean field with lattice symmetry. It is expected[1] that vacancy-interstitial (VI) fluctuations play an important role in supersolidity. In the JN function, VI pairs are bound, just as electron-hole pairs are in an electronic insulator. In the metallic function, vacancies will be locally attracted to interstitials but they are not bound as pairs; as a result the Ψ_M has both BEC and NCRI. Calculations on a similar quantum solid[18] have found that Ψ_M has a higher energy than Ψ_{JN} ; it costs energy to create unbound VI fluctuations. We note that calculations[20] and experiments[21] on solid ^4He suggest that there are no unbound vacancies or interstitials at low temperatures.

Another trial function, the “shadow” wave function[22] (SWF) has been introduced to allow a translationally invariant trial function to have the correct solid order. In this case, a single atom coordinate in the Jastrow trial function is replaced by a composite object, a linear polymer, with the inter- and intra- correlation factors being variationally determined. Typically the polymer is in fact a dimer, with the two ends separated by a distance on the order of the inter-atomic spacing. One of the ends (the shadow) is integrated over to get the trial function. The great advantage of the SWF is that the crystal order is spontaneously generated, and one gets accurate properties after optimizing the wave function parameters. As more monomers are added to the trial function, the functional form is equivalent to the Feynman-Kacs path integral expression for the ground state wave function[23]. Though it has not been analytically shown, simple argu-

ments involving displacing the polymer, make it is plausible that the shadow trial function will have always have BEC, as long as the inter-atomic correlation factors are short-ranged. This has been verified in recent numerical calculations by Galli et al.[24]

Thus, we have a dilemma, not uncommon with arguments based on variational wavefunctions: one can have several satisfactory trial functions, all of them capable of good descriptions of solid helium, but some are BEC and some are not BEC. All of the usual properties one uses to test the quality of the trial function, *e. g.* the energy, the Debye-Waller factor, the pair correlation function, etc. are diagonal in coordinate space (expectations of $|\Psi(R)|^2$); hence they are unreliable measures of how accurate the off-diagonal elements needed in Eq. (1) are. As we have mentioned above, it is not possible to state a general theorem covering whether all quantum crystals must or cannot have BEC, since the Jastrow trial functions can be used to define a Hamiltonian which is supersolid. We need a more direct, reliable method to decide whether the ground state of solid ^4He is BEC and, hence, which class of trial function, metal or insulator, is appropriate. Calculating $n(\Delta\mathbf{r})$ in solid bulk ^4He allows us to establish how good these trial functions are for computing BEC in quantum crystals.

To provide an unbiased answer, we use path integral Monte Carlo (PIMC)[25], a numerical method that calculates integrals over the many body density matrix. It is ideally suited for this calculation since it can be done at finite temperature (under conditions where an experimental signature of NCRI has been seen), is, in principle, exact, and has been validated on many properties[25] of liquid and solid ^4He . Most importantly, it is independent of a trial wave function bias or any assumption of lattice. Only the He-He potential enters: a semi-empirical form[26] is known to be accurate; in any case, experimental results suggest that supersolid behavior is a robust phenomenon insensitive to fine details of the interaction.

In PIMC, based on Feynman’s description of superfluid helium, one maps the quantum system onto a classical system of “polymers” which can permute onto one another. The value of $n(|\Delta\mathbf{r}|)$ is mapped to the end-to-end distribution of an open “polymer” in a sea of closed “polymers” representing the other helium atoms. If the system has ODLRO, the two ends will become separated from each other; the condensate fraction is the value of $n(\Delta\mathbf{r})$ at large r . On the other hand, if there is no ODLRO, the two ends will remain localized relative to each other and $n(\Delta\mathbf{r})$ will approach 0 at large $|\Delta\mathbf{r}|$. Superfluidity (*i.e.* NCRI), on the other hand, is represented in PIMC as long cyclic permutation of paths, for example, as a complete row of atoms shifting down one element. One can immediately see the difference between ODLRO and NCRI in a PIMC solid calculation. Suppose we have N atoms and N lattice sites, so each site is, on the average, singly occupied. To perform the ODLRO

calculation, one of the atoms paths is opened up, so now there will be $N - 1$ closed polymers and 2 open ends; clearly one of the N sites must be doubly occupied if we count the open ends as distinct atoms. For NCRI one can simultaneously shift all atoms to their new permuted sites; there need not be any double occupation, but such a simultaneous shift is unlikely for many atoms.

Techniques[25] developed and tested[28] on superfluid ^4He are used to calculate $n(\Delta\mathbf{r})$ efficiently, although new moves are used to ensure ergodicity in the solid. If one simply cuts open one polymer, one rarely finds large values of $\Delta\mathbf{r}$. To guarantee sufficient statistics on the end-end distance, we multiply the path integral density by a series of importance sampling functions having the effect of forcing apart the ends. A series of importance functions are used to get good statistics for a specific range of magnitudes of $|\Delta\mathbf{r}|$. The separate end-to-end distributions are then normalized with respect to each other by minimizing the χ^2 difference between them and using the definition $n(0) = 1$. We perform calculations of $n(\Delta\mathbf{r})$ both by performing an angular average over $\Delta\mathbf{r}$ as well as fixing $\Delta\mathbf{r}$ in the basal plane, specifically in the x direction. Calculations of $n(\Delta\mathbf{r})$ can much more easily access the full path space than calculations of the superfluid density because the ends of the open polymer can move more freely.

Calculations have been performed in a nearly cubic box of 180 particles as well as a series of longer rectangular boxes extended in the basal plane ranging from 48 to 144 atoms. For the longer box, we altered our importance function to force the open ends apart only in the x direction (in the nearest neighbor direction of the basal plane). We examine values of $n(\Delta\mathbf{r})$ for a range of temperatures from $0.1K$ to $2K$ in a (periodic) rectangular box of size $\sim 18.5 \text{ \AA}$ corresponding to a density, $0.0286 \text{ \AA}^{-3}$ near the experimental melting pressure of 25.3 bars. The random walk is started with the atoms in a perfect hcp crystalline lattice but no constraints are placed on how the atoms can arrange themselves. Since the box is chosen to be commensurate with a perfect hcp crystal, vacancies and interstitials are expected not to exist in the system though they can form by a fluctuation. We have verified that the results are independent of temperatures below $2.0K$.

The results of these calculations for $T = 0.5K$ are shown in figure 1. Note the curvature at small $\Delta\mathbf{r}$ is determined by the kinetic energy estimated at $24.0 K/\text{atom}$. For both the spherical average value and the value in the basal plane $n(|\Delta\mathbf{r}|)$ decays exponentially for $|\Delta\mathbf{r}| > 3\text{\AA}$: we find no indication of ODLRO (BEC) in bulk solid ^4He . In fact the rate of decrease for $n(x)$, the SPDM in the x -direction, is in agreement with the frequencies for straight line winding exchange[5], shown as the solid line. We find oscillations in the computed $n(x)$ reflecting the underlying crystal lattice. The angular averaged $n(r)$ shows less structure than that in the x -direction. Though it is pos-

sible that $n(x)$ will plateau for a longer distance scale, we see no indication of this in the results, nor can we think of a physical mechanism that would be responsible[29]. A system with ODLRO will be manifested when one end of the open polymer loses knowledge about the other end of the open polymer. But we find that pulling apart the two ends takes “work” per unit length, no matter how far apart the two ends are. This happens because additional atoms are displaced from their lattice sites.

The results of our PIMC calculations are in quantitative agreement with variational Monte Carlo calculations using a shadow wave function[24] for distances less than $9 \text{ \AA}$; they conclude that BEC does exist in the ground state of ^4He with a very small condensate fraction, $n_0 = 5 \times 10^{-6}$ at the melting density. However, we find that $n(\Delta\mathbf{r})$ continues to decrease beyond $9 \text{ \AA}$.

We believe the reason for the discrepancy with PIMC is that the shadow wave function is a technique that improves upon the Jastrow wave function, in fact it is remarkably accurate up to the second neighbor distance. Nonetheless the wavefunction is still built from short range correlations. This leads inevitably to a non-zero superfluid fraction. We also expect the SWF will give a non-zero but very small vacancy concentration[30]. PIMC does not make a variational ansatz, with the sole assumption that the results at low temperature smoothly approach the ground state values.

Among the proposed quantum solid wave functions, the symmetry-breaking Nosanow-Jastrow wavefunction, gives results in agreement with PIMC, validating the “insulator” representation of solid helium. (Further details about this picture are in ref. [15]). The implication of our calculation and the Reatto-Chester theorem is that a quantum solid trial function must include long-range correlations. This is most simply done by putting in localized functions about lattice sites, reflecting the broken symmetry. We have only done PIMC calculations at the melting density but we do not expect different behavior at higher density, since difficulty of exchange grows rapidly with density. The result reported here, together with the finding of zero superfluid density reported in ref. [5] suggest that the mechanism for the measurements of Kim and Chan[4] involves more than equilibrium properties of a perfect ^4He crystal.

(After the calculation reported here was completed, we have learned of a similar PIMC calculation [32]. The overall conclusions are similar, however their Monte Carlo algorithm and results for $n(\Delta\mathbf{r})$ differ in important ways.)

This work was supported by NSF under grant no. DMR-03 25939 ITR and the fundamental physics program at NASA (NAG-8-1760). Computer time has been provided by NCSA and the Materials Computation Center and the Frederick Seitz Materials Research Laboratory (U.S. DOE DEFG02-91ER45439) at the University of Illinois Urbana-Champaign. We acknowledge useful




FIG. 1: (color online) The single particle density matrix as a function of distance (in Å), estimated with PIMC, in hcp ^4He at a density 286nm^{-3} and a temperature of 0.5K. The red curve is the spherically averaged $n(r)$ while the blue curve is $n(x)$. The upper (green) curve is a variational MC calculation using the shadow wave function ref. [24]. The straight line (black) has a slope determined from the exchange calculations of ref. [5].

communications with D.Galli, L. Reatto, W. Saslow, N. Trivedi and G. V. Chester.

-
- [1] A. F. Andreev and I. M. Lifshitz, Sov. Phys. JETP **29**, 1107 (1969).
 - [2] A. J. Leggett, Phys. Rev. Lett. **25**, 1543 (1970).
 - [3] G. V. Chester, Phys. Rev. A **2**, 256 (1970).
 - [4] E. Kim and M. H. Chan, Science **305**, 1941 (2004).
 - [5] D. M. Ceperley and B. Bernu, Phys. Rev. Letts. **93**, 155303 (2004).
 - [6] O. Penrose and L. Onsager, Phys. Rev. **104**, 576 (1956). Letts. **94**, 155302 (2005).
 - [7] J. A. Hodgdon and F. H. Stillinger, J. Stat. Phys. **78**, 117 (1995).
 - [8] Phys. Rev. **183**, 334 (1969).
 - [9] To prove that there exist supersolids, assume that the exact bosonic wavefunction equals the potential of a known classical crystal, e.g. a Lennard-Jones crystal[7]. Then define the Hamiltonian of the quantum system to have this as an exact ground state; it must be the ground state since it is nodeless. The quantum potential as defined is not a pair potential; it has a very strong 3-body potential.

The crystal will also have a non-zero density of vacancies since the equivalent classical system does.

- [10] N. Prokof'ev and B. Svistunov, Phys. Rev. Lett. **94**, 155302 (2005).
- [11] J. P. Hansen, D. Levesque, Phys. Rev. **165**, 293 (1968); J. P. Hansen and E. L. Pollock, Phys. Rev. A **5**, 2651 (1972).
- [12] C. A. Burns and E. D. Isaacs Phys. Rev. **B 55**, 5767 (1997).
- [13] L. H. Nosanow, Phys. Rev. Lett. **13**, 270 (1964)
- [14] M. Schwartz, Phys. Rev. B **12**, 3725 (1975).
- [15] P. W. Anderson, *Basic Notions of Condensed Matter Physics*, Westview Press, 1984, see chapter 4.
- [16] Although it is true that the exact wave function for a finite system will have translation invariance (TI), a trial function lacking TI will be closer to the thermodynamic limit since if one averages over the center-of-mass, that lowers the energy by order $(1/N)$, but such motion is unphysical in the limit of large N .
- [17] D. N. Lowy and C.-W. Woo, Phys. Rev. B **13**, 3790 (1976).
- [18] D. Ceperley, G. V. Chester, M. H. and Kalos, Phys. Rev. B **17**, 1070 (1978).
- [19] H. Zhai and Y-S Wu, J. of Stat. Mech. in press (2005); cond-mat/0501309.
- [20] B. Chaudhuri, F. Pederiva, and G. V. Chester, Phys. Rev. **B 60**, 3271 (1999).
- [21] B. A. Fraas, P. R. Granfors, and R. O. Simmons, Phys. Rev. **bf B 39**, 124 (1989).
- [22] S. Vitiello, K. Runge, and M. H. Kalos Phys. Rev. Lett. **60**, 1970 (1988).
- [23] D. E. Galli and L. Reatto, Mol. Phys. **101** 1697 (2003).
- [24] D. Galli, M. Rossi and L. Reatto, Phys. Rev. B **71**, 140506 (2005).
- [25] D. M. Ceperley, Rev. Mod. Phys. **67**, 279 (1995).
- [26] R. A. Aziz, A. R. Janzen and M. R. Moldover, Phys. Rev. Lett. **74**, 1586 (1995).
- [27] D. M. Ceperley and E. L. Pollock, Can. J. Phys. **65**, 1416 (1987).
- [28] We used pair density matrices and found an imaginary-time step of $0.0125K^{-1}$ was sufficient for convergence.
- [29] The observed transition temperatures rule out very small condensate fractions since one expects that if ODLRO is due to BEC of point defects then $n_0 \sim \hbar^3(mk_B T_c)^{3/2} \sim 0.01$.
- [30] The concentration of vacancies is given directly by the square of the trial wavefunction, $|\Psi|^2$ in the thermodynamic limit, e.g. considered as a classical Boltzmann distribution[7]. For approximate trial functions, this will give a different answer than a variational estimate of the vacancy energy from the Hamiltonian: $E_v(N \pm 1, N) - E_v(N, N)$ as was done in ref. [31]. Here, $E_{var}(N, N_s)$ is the variational energy of N atoms and N_s lattice sites at constant atomic density.)
- [31] F. Pederiva, G. V. Chester, S. Fantoni and L. Reatto, Phys. Rev. B **56**, 5909 (1997).
- [32] M. Boninsegni, N. Prokof'ev, and B. Svistunov, cond-mat/0512103.